

## Laser based photoelectron circular dichroism: from femtoseconds to nanoseconds

Exploiting an electric dipole effect in ionization [1], photoelectron circular dichroism (PECD) is a highly sensitive enantioselective spectroscopy for studying chiral molecules in the gas phase using either single-photon ionization [2] or multiphoton ionization [3]. In the latter case resonance enhanced multiphoton ionization (REMPI) gives access to neutral electronic excited states. The PECD sensitivity opens the door to study control of the coupled electron and nuclear motion in enantiomers. A prerequisite is a detailed understanding of PECD in REMPI schemes. In this contribution I will report on our recent experiments devoted to unravel different aspects of this effect on the fenchone prototype by addressing the range from impulsive excitation on the femtosecond time scale to highly vibrational state selective excitation with the help of high resolution nanosecond laser techniques [4]. Subcycle effects in bichromatic fields are shortly addressed as well [5].

[1] B. Ritchie, *Phys. Rev. A* 1976, 13, 1411–1415.

[2] N. Böwering, T. Lischke, B. Schmidtke, N. Müller, T. Khalil, U. Heinzmann, *Phys. Rev. Lett.* 2001, 86, 1187–1190.

[3] C. Lux, M. Wollenhaupt, T. Bolze, Q. Liang, J. Köhler, C. Sarpe, T. Baumert, *Angew. Chem. Int. Ed.* 2012, 51, 5001–5005.

[4] A. Kastner, G. Koumarianou, P. Glodic, P. C. Samartzis, N. Ladda, S. T. Ranecky, T. Ring, S. Vasudevan, C. Witte, H. Braun, H.-G. Lee, A. Senftleben, R. Berger, G. Barratt Park, T. Schäfer, T. Baumert, *Phys. Chem. Chem. Phys.*, 2020, 22, 7404.

[5] P. V. Demekhin, A. N. Artemyev, A. Kastner, T. Baumert, *Phys. Rev. Lett.*, 2018, 121, 253201.